

Tutorial 5 : One-point compactification

①
updated 4/9

The two "nicest" classes of spaces are metrisable spaces, and compact Hausdorff spaces. After that, the next best kind of space are subspaces of these kinds of spaces : subspaces of metrisable spaces are metrisable but subspaces of compact spaces need not be compact, so in this latter case we obtain a new and interesting class of spaces : the locally compact Hausdorff spaces. In the second half of this course we will prove many theorems about this class of spaces. In this tutorial we develop the theory of locally compact spaces, as an extended exercise in the basic properties of compactness itself.

Defⁿ A topological space X is locally compact if for every $x \in X$ there exists an open set U and compact set K with $x \in U \subseteq K$.

Clearly any compact space is locally compact.

Example (i) \mathbb{R}^n is locally compact (but not compact) since any $x \in \mathbb{R}^n$ lies in some $(a_1, b_1) \times \dots \times (a_n, b_n)$ which is contained in the compact set $[a_1, b_1] \times \dots \times [a_n, b_n]$.

(ii) \mathbb{Q} is not locally compact (so subspaces of locally compact spaces need not be locally compact).

Defⁿ A topological space X is Hausdorff if for any pair $x, y \in X$ of distinct points there exist open $U, V \subseteq X$ with $x \in U, y \in V$ and $U \cap V = \emptyset$. (developed in L11)

Remark (i) In a Hausdorff space points are closed.

(ii) Any subspace of a Hausdorff space is Hausdorff.

(iii) A compact subspace of a Hausdorff space is closed.


Q1 Any closed subspace of a locally compact space is locally compact.

Theorem If X is locally compact Hausdorff then it is homeomorphic to a subspace of a compact Hausdorff space. (e.g. $X = \mathbb{R}$)

We define for locally compact Hausdorff X a space \tilde{X} that we call the one-point compactification, which is compact Hausdorff and has X as a subspace.

Defⁿ As a set $\tilde{X} = X \sqcup \{\infty\}$. A subset $U \subseteq \tilde{X}$ which does not contain ∞ is open iff. it is open in X , and $U \subseteq \tilde{X}$ which contains ∞ is open iff. there is $K \subseteq X$ compact with $U = K^c \sqcup \{\infty\}$.

Example $X = \mathbb{R}^2$ is locally compact



The set $K^c \sqcup \{\infty\}$ is an open neighborhood of ∞ in \tilde{X} . By Heine-Borel a compact subset $K \subseteq \mathbb{R}^2$ must be closed and bounded, so this picture is actually generic, i.e. it describes all open neighborhoods of ∞ .

[Q2] Prove this is a topology on \tilde{X} .

[Q3] Prove \tilde{X} is compact and Hausdorff.

[Q4] Prove the canonical map $X \rightarrow \tilde{X}$ induces a homeomorphism between X and a subspace of \tilde{X} (hence proving the theorem).

This space \tilde{X} is called the one-point compactification of X . It is not a very smart construction, the Stone-Cech compactification is better, but the one-point compactification has its uses.

[Q4] Give a characterisation of the set of continuous functions $f: X \rightarrow \mathbb{R}$ which extend to \tilde{X} , that is, for which $F: \tilde{X} \rightarrow \mathbb{R}$ continuous exists making the diagram below commute:

$$\begin{array}{ccc} X & \xrightarrow{\quad} & \tilde{X} \\ & \searrow f & \downarrow F \\ & & \mathbb{R} \end{array}$$

This class of functions will be necessary for defining the Hilbert spaces $L^2(\mathbb{R}^d)$ of square-integrable functions on Euclidean space.